

兔抗 DOK2 (Phospho-Tyr299)多克隆抗体

中文名称:	兔抗 DOK2 (Phospho-Tyr299)多克隆抗体
英文名称:	Anti-DOK2 (Phospho-Tyr299) rabbit polyclonal antibody
别名:	p56DOK; p56dok-2
相关类别:	一抗
储存:	冷冻 (-20℃) 避光
宿主:	Rabbit
抗原:	DOK2 (Phospho-Tyr299)
反应种属:	Human
标记物:	Unconjugate
克隆类型:	rabbit polyclonal

技术规格

Background:	DOK proteins are enzymatically inert adaptor or scaffolding proteins. They provide a docking platform for the assembly of multimolecular signaling complexes. DOK2 may modulate the cellular proliferation induced by IL-4, as well as IL-2 and IL-3. May be involved in modulating Bcr-Abl signaling. Attenuates EGF-stimulated MAP kinase activation
Applications:	WB, IHC, IF
Name of antibody:	DOK2 (Phospho-Tyr299)
Immunogen:	Synthetic peptide of human DOK2 (Phospho-Tyr299)
Full name:	docking protein 2, 56kDa (Phospho-Tyr299)
Synonyms :	p56DOK; p56dok-2
SwissProt:	O60496
IHC positive control:	Human breast carcinoma

IHC Recommend dilution:	50-100
WB Predicted band size:	56 kDa
WB Positive control:	Jurkat cells treated with insulin
WB Recommended dilution:	500-1000
IF Positive control:	Hela cells
IF Recommended dilution	100-200



